




PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁷ : C12N 15/54, 9/12, C12Q 1/68</p>	<p>A1</p>	<p>(11) International Publication Number: WO 00/53772 (43) International Publication Date: 14 September 2000 (14.09.00)</p>
<p>(21) International Application Number: PCT/EP00/01917 (22) International Filing Date: 4 March 2000 (04.03.00) (30) Priority Data: 99104539.4 6 March 1999 (06.03.99) EP 99117245.3 2 September 1999 (02.09.99) EP (71) Applicant (for all designated States except US): ROCHE DIAGNOSTICS GMBH [DE/DE]; D-68298 Mannheim (DE). (72) Inventors; and (75) Inventors/Applicants (for US only): KAEHLER, Markus [DE/DE]; Klosterhof 8, D-25524 Itzehoe (DE). FREY, Bruno [DE/DE]; Hochfeldstrasse 50, D-82377 Penzberg (DE). SOBEK, Harald [DE/DE]; Birkenstrasse 29, D-82377 Penzberg (DE). ANTRANIKIAN, Garabed [DE/DE]; Schafkovengrund 3, D-21218 Sevetal-Hittfeld (DE). (74) Common Representative: ROCHE DIAGNOSTICS GMBH; Patent Department, D-68298 Mannheim (DE).</p>	<p>(81) Designated States: JP, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published With international search report.</p>	

(54) Title: DNA POLYMERASE FROM PYROBACULUM ISLANDICUM



(57) Abstract

The invention is related to a DNA polymerase obtainable from *Pyrobaculum islandicum* whereas the six conserved motifs indicative of family B DNA polymerases were contained in the *Pyrobaculum islandicum* DNA polymerase sequence and whereas this polymerase exhibits 3'-5'-exonuclease activity.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

DNA Polymerase from *Pyrobaculum islandicum*

The invention is related to a DNA polymerase obtainable from *Pyrobaculum islandicum* whereas the six conserved motifs indicative of family B DNA polymerases were contained in the *Pyrobaculum islandicum* DNA polymerase sequence and whereas this polymerase exhibits 3'-5'-exonuclease activity.

Archaea constitute the third domain of life, distinguishable from the domains Eukarya and Bacteria (Woese and Fox 1977). During the last two decades, many Archaea have been isolated and characterized. Hyperthermophilic Archaea, which are the deepest and shortest branches of the phylogenetic tree, can grow at temperatures up to 110°C (Stetter 1996). Thus, these organisms are excellent candidates to investigate their thermostable enzymes. While our knowledge of DNA replication in Eukarya and Bacteria is quite advanced (Kornberg and Baker 1992), only limited information on the replication of DNA in hyperthermophilic Archaea is available. It is still an open question how genomes can be maintained and duplicated at temperatures near or above 100°C.

More than 50 DNA polymerase genes from Eukarya, Bacteria and Archaea have been cloned and sequenced. Deduced from their amino acids sequences, these enzymes can be classified into four groups, represented by *Escherichia coli* DNA polymerase I (family A), DNA polymerase II (family B), DNA polymerase III (family C), and eucaryotic DNA polymerase β (family X) (Braithwaite and Ito 1993). The majority of archaeal DNA polymerases described so far belongs to the family B. A number of euryarchaeal DNA polymerase genes has been cloned and sequenced. The thermostable enzymes from several *Thermococcus* and *Pyrococcus* strains have been successfully expressed, characterized and can be obtained commercially (Perler et al. 1996). Relatively little information is available on DNA polymerases from hyperthermophilic crenarchaeota. Uemori et al. (1995) described the cloning and characterization of two family B DNA polymerases from the hyperthermophilic crenarchaeon *Pyrodicticum occultum*. Despite of the fact that three family B DNA polymerase genes have been detected in *Sulfolobus solfataricus* (Prangishvili et al. 1993; Datukishvili et al. 1996; Edgell et al. 1997), only one of these genes has been successfully expressed (Pisani et al. 1992).

DNA polymerases from hyperthermophiles play a major role in a variety of molecular biological applications, e.g. DNA amplification, sequencing, labelling or reverse transcription. The develop-

ment of polymerase chain reaction (PCR; Mullis et al. 1986; Erlich et al. 1988) was one of the most important contributions to biotechnology in the last decade. The DNA polymerase I from the bacterium *Thermus aquaticus*, called *Taq* polymerase, was the first thermostable DNA polymerase applied in PCR. Due to the absence of a 3'-5' exonuclease activity, this enzyme is unable to excise mismatches and as a result, the base insertion fidelity is low (Longley et al. 1990; Keohavong and Thilly 1989). Furthermore, research has indicated that *Taq* polymerase has a thermal stability of not more than few minutes at 100°C. Several euryarchaeal DNA polymerases with much higher thermostability and an associated 3'-5' exonuclease activity, e.g. *Pfu* pol (Lundberg et al. 1991) from *Pyrococcus furiosus* or Vent pol (Cariello et al. 1991) from *Thermococcus litoralis*, have been described and are commercially available. However, *Taq* polymerase was not replaced by these polymerases mainly due to their low extension rates. Therefore, new DNA polymerases with enhanced thermostability, high base insertion fidelity and sufficient extension rates are still needed. Hyperthermophilic crenarchaeota could be a promising source for thermostable DNA polymerases with new properties. Additionally, the production of large quantities of recombinant enzymes would be of great value.

The anaerobic hyperthermophilic Archaeon *Pyrobaculum islandicum*, which belongs to the crenarchaeota, was isolated from an Icelandic hot spring (Huber et al. 1987). It was not possible to isolate polymerases from the native strain without undue experimentation because this strain is difficult to culture. Furthermore, it was not possible to amplify fragments of the B DNA polymerase gene of *Pyrobaculum islandicum* using conventional primers.

PCR Primer Screening

In order to amplify fragments of the *Pyrobaculum islandicum* DNA polymerase, we performed a primer screening with various kind of primers. Archaeal family B DNA polymerase are difficult to align due to short, highly conserved domains separated by long stretches of low or no amino acid conservations. It seems to be particularly difficult to clone a new family B DNA polymerase gene from an archaeon which shows phylogenetically large distances to archaea from which polymerase sequences have been identified.

Pyrobaculum islandicum belongs to the crenarchaeal order Thermoproteales. When we started our investigations, only a few crenarchaeal polymerase genes from *Pyrodictum* and *Sulfolobus* species, and some polymerase gene sequences from the euryarchaeal *Pyrococcus* and *Thermococcus* species have been identified.

The PCR primer screening (using *Pyrobaculum islandicum* DNA as template) was started with the combination of several primers deduced from conserved *Thermococcus* and *Pyrococcus* polymerase sequences:

forward primers

tcg1:	5'-CCGCAAATGATACTGGACACTGATTACATAAC-3'	SEQ ID No. 5
pcw1:	5'-CCGCAAATGATTTTAGATGTGGATTACATAAC-3'	SEQ ID No. 6
c1435:	5'-GAGCCAGAAAARGGKTTGTGGGAAAA-3'	SEQ ID No. 7

reverse primers

tcg2:	5'-CCGCAATTAGGATTTTTTAATGTTAAGCCAGGAAG-3'	SEQ ID No. 8
pcw2:	5'-CCGCAATTACCTCTTGAGCCATGCATCTAAGCCGG-3'	SEQ ID No. 9
c1190:	5'-TCATAMGTTGCCYTWGCATCTTCCAT-3'	SEQ ID No. 10

The primer pair tcg1/c1190 was successfully used before to amplify a *Thermococcus* polymerase gene fragment (Niehaus et al. 1997).

In contrast to the *Thermococcus* polymerase gene however, it was not possible to amplify any PCR products from *Pyrobaculum islandicum* showing identities to family B DNA polymerases.

Uemori et al. (1995) also used degenerated primers, deduced from highly conserved regions of family B DNA polymerases genes:

forward, 5'-(TA)(CG)(CG)(CT)T(CG)TAC CC(CG) (TA)(CG) ATC AT-3'
reverse, 5'-TC(TCAG) GT(AG) TC(TCAG) CC(AG) TA(AG) AT(TCAG) AC-3

A PCR with these primers and *Pyrobaculum islandicum* DNA as template was also not successful.

Therefore, it was necessary to design new degenerated primers. Additionally, it can be assumed that the use of degenerated primers, including deoxyinosit (I), may be a great advantage in amplifying new archaeal genes, because primers with deoxyinosit accept several degenerated triplets.

The two available family B DNA polymerase genes from *Pyrodictium occultum* (Uemori et al. 1995; ass. no. D38574 and D38573) and one polymerase gene from *Sulfolobus solfataricus* (ass. no. Y08257) was chosen for an alignment in order to synthesize suitable primers for a primer screening.

Due to the degenerated triplett code, it was necessary to design degenerated primers, using deoxyninosin (I). The primers were deduced from highly conserved regions of family B DNA polymerases.

forward primers

f1:	5'-CTG ACC TT(CT) AA(CT) GGI GA(CT)-3'	SEQ ID No. 11
f2:	5'-CAT ATI GAC (CT)TI TA(CT) AAI TT-3'	SEQ ID No. 12
f3:	5'-GTI (CT)TI GAT TT(CT) GCI (AT)(GC)I (CT)TI TA(CT) CC-3'	SEQ ID No. 13
f4:	5'-ATI AAG GGI AA(AG) AAG TAT-3'	SEQ ID No. 14

reverse primers

r1a:	5'-(CT)TC IGG IGT (AG)TT IC(GT) (CT)TT (CT)TT IGC-3'	SEQ ID No. 15
r1b:	5'-GAA IAG GTC IGT ATC ICC AT-3'	SEQ ID No. 16
r2a:	5'-GAA IAG IGA (AG)TC IGT (AG) TC ICC (AG)T-3'	SEQ ID No. 17
r2b:	5'-GAA IAG IGA GTC IGT ATC ICC AT-3'	SEQ ID No. 18

A multiple primer screening was performed with combinations of each forward with each reverse primer. A low annealing temperature of 48°C was used.

Only one PCR with the forward primer f3 and the reverse primer r2b resulted in the amplification of a PCR product, showing identities to archaeal family B DNA polymerases. This fragment was used as a probe to identify the complete polymerase gene, as described in Example 2. This result indicated that it is very difficult to design appropriate primers for the amplification of new crenarchaeal DNA polymerase gene fragments.

The subject of the present invention comprises the cloning procedure and sequence of a DNA polymerase gene from *Pyrobaculum islandicum*. The invention comprises a gene with SEQ ID No. 3. The subject of the present invention also comprises a B DNA polymerase from *Pyrobaculum islandicum*. The invention comprises a polymerase with a sequence according to SEQ ID No. 4. Furthermore, the expression, purification and characterization of the recombinant polymerase from *P. islandicum* is disclosed.

Cloning and nucleotide sequencing of the DNA polymerase gene from *P. islandicum*

Based on the most conserved amino acid sequences found in archaeal α -like DNA polymerases, two degenerated primers according to SEQ ID No. 1 and SEQ ID No. 2 were designed. The PCR was performed with chromosomal DNA from *P. islandicum*, and we were able to obtain an amplicate

of 445 bp. The PCR product was sequenced, and the sequence showed the expected homology to known archaeal polymerase regions in a BLAST alignment. As an important result of the present invention, known archaeal DNA polymerase sequences can be successfully used to design novel degenerated primers in order to find new DNA polymerase genes from hyperthermophilic archaea. Therefore, the cloning procedure using degenerated primers according to SEQ ID No. 1 and SEQ ID No. 2 is also subject of the present invention. The PCR product was then used as a probe to obtain the whole DNA polymerase gene. In southern blot hybridizations, *P. islandicum* DNA fragments in a size of 6 to 7.5 kbp, digested with *SacI*, were chosen to be cloned in a cosmid vector system. An open reading frame consisting of 2356 bp and coding for a protein with 784 amino acids (Fig. 1) was detected on an insert of a cosmid clone, which was identified by colony filter hybridization. The open reading frame started with GTG, coding for valin. It has been shown that some archaeal proteins (approx. 25 %) appear to start at GTG codons (Dennis 1997). For example, Schleper et al. (1997; 1998) described several open reading frames in the crenarchaeote *Cenarchaeum symbiosum*, including that of a family B DNA polymerase gene, starting with GTG instead of ATG. In order to examine the putative transcription and translation start region of the DNA polymerase gene from *P. islandicum*, a 5'RACE PCR was performed and the amplified cDNA was sequenced. The sequence of the mRNA started one or two nucleotides upstream of the determined start codon (data not shown).

The DNA polymerase sequence was aligned with those of all archaeal DNA polymerases available in public databases. The six conserved motifs indicative of family B DNA polymerases (Braithwaite and Ito 1993) were identified in the *P. islandicum* sequence, including those residues that are invariant in all DNA polymerases and have been shown to be functionally important. Also, the three 3'-5' exonuclease motifs (Blanco et al. 1991) were localized in the sequence. In a multiple alignment performed with the ClustalW software program (Higgins, EMBL, Heidelberg, Germany), the DNA polymerase sequence showed highest identities to one of the two family B DNA polymerases from *Pyrodictium occultum* (polB; accession. no. D38574), a family B DNA polymerase from *Archaeoglobus fulgidus* (accession no. AE001070) and the *Pfu* polymerase (accession no. D12983), sharing 51%, 37% and 32% identical residues, respectively.

Expression and purification of *P. islandicum* DNA polymerase

After determining the sequence, we cloned the gene into vector pASK-IBA 3 and expressed the polymerase as a fusion protein with a C-terminal Streptactin-binding oligopeptide. The recombinant DNA polymerase was purified to near homogeneity by the combination of a heat denaturation step and an affinity chromatography on Streptactin. Starting with 1000 ml of *E. coli* BL21

culture, we purified 1 mg of recombinant DNA polymerase with a specific activity of approx. 2000 U/mg preparations of recombinant DNA polymerase. Attempts to purify the polymerase without addition of protease inhibitors to the buffer W resulted in the appearance of many small contaminants as judged by SDS-PAGE (data not shown). The purity of the DNA polymerase in each fraction were checked by SDS-PAGE (Fig. 2). The fusion protein was copurified with a small amount of a ca. 70 kDa protein as a contaminant. The activity gel illustrated that the purified protein of approx. 90 kDa in the Streptactin preparation possesses DNA polymerase activity (Fig.3). A small activity was also detectable at approx. 70 kDa, indicating that the copurified protein was a proteolytic degradation product of the recombinant polymerase. Additionally, in crude extract from *P. islandicum*, we were able to detect corresponding DNA polymerase activity in a protein band of approx. 90 kDa.

Subjects of the present invention are also mutants of the inventive B DNA polymerase and DNA sequencing encoding these mutants. These mutants may exhibit e.g. reduced proofreading activity according to mutations in the highly preserving amino acid regions for this exonuclease activity (ExoI, ExoII and ExoIII) e.g. ExoI region contains an X1DX2EX3 motif, and the amino acids D (aspartic acid) and E (glutamic acid) are known to be essential for the exonuclease activity for B type polymerases.

Biochemical properties

To determine the biochemical properties of the purified recombinant DNA polymerase, we carried out the non-radioactive assay described in Materials and Methods under different reaction conditions.

The optimal conditions for DNA polymerase activity were tested at 75°C in the presence of activated calf thymus DNA as a template. As shown in Fig. 4, the polymerase was most active at pH 7.3. The optimal temperature for activity was not detectable because activated DNA was not stable above 80°C. The enzyme required an extremely low concentration of divalent cations for activity. Concentrations above the optimum of 0.5 mM MgCl₂ inhibited the polymerase (Fig. 5). No activity was detected in the absence of divalent cations. As illustrated in Fig. 6, monovalent potassium ions strongly inhibited the DNA polymerase. In the presence of 100 mM KCl, only 7% of the residual activity was detected.

To determine the thermostability of the DNA polymerase, a preincubation of the enzyme in 50 mM Tris-HCl pH 7.2 was performed at 90 and 100°C, following by an incubation at 75°C under stan-

standard assay conditions. Half-life times of 30-40 min at 100°C and >10 h at 90°C have been detected (Fig. 7).

The influence of representative inhibitors on the DNA polymerase activity was investigated (Tab. 1). The sulfhydryl blocking reagent N-Ethylmaleimide (NEM) strongly inhibited the DNA polymerase activity. In the presence of 1 mM NEM, 40% residual activity was detected. A total loss of activity was detected after a preincubation with 5 mM NEM. The sensitivity of the enzyme to the tetracyclic diterpenoid aphidicolin, which competes with each dNTP for binding to DNA polymerase, was also investigated. The DNA polymerase was inhibited by a comparable amount of aphidicolin that inhibits polB from *Pyrodicticum occultum* and *Pfu* polymerase (Uemori et al. 1995). In the presence of 500 and 2000 µM aphidicolin, we detected 55 and 21% residual activity for the *P. islandicum* DNA polymerase. A very low sensitivity of the polymerase to ddGTP was detected. No inhibition was measured at a ddGTP/dGTP ratio of 2, and at a ddGTP/dGTP ratio of 10, approx. 80% of residual activity could be detected.

In order to determine associated exonuclease activity in the DNA polymerase from *P. islandicum*, we first incubated *Bam*HI digested pUC 19 DNA with the purified enzyme in the absence of dNTP. A significant degradation after 2 h at 75°C was detected on an agarose gel (data not shown). No degradation was measured at 37°C, indicating that no exonuclease contamination from the *E. coli* BL21 cells was present in the enzyme preparation after heat-treatment and purification on Strep-tactin Sepharose.

The exonuclease activity was quantified in a standard assay, based on the release of ³H-labeled nucleotides from a sonicated calf thymus DNA template. The release of nucleotides after an incubation of the template with 1.5 U purified polymerase at 65°C was comparable to that caused by other archaeal proofreading DNA polymerases. 104 cpm/10 min were measured after 4 h incubation. In comparison, 15 U (10-fold higher concentration) of *Taq* polymerase caused a cpm/10 min of 150-250, and the same amount of proofreading polymerases such as *Pfu* polymerase caused 1000-3000 cpm/min. No release of nucleotides was detected after incubations at 37°C, indicating that no *E. coli* exonuclease activity interferes with the test. The detected exonuclease activity of the *P. islandicum* DNA polymerase corresponds to the three 3'-5' exonuclease domains identified within the deduced DNA polymerase sequence (Blanco et al. 1991), indicating that this kind of exonuclease activity and not a 5'-3' exonuclease activity was measured.

Additionally to the above described exonuclease activity assay 3'-5' exonuclease activity was tested on single strands as described below. To investigate if a 3'-5' exonuclease activity on single- and

double-stranded DNA could be detected, we used a 5'-DIG-labeled primer and a primer/template complex. 3'-5' exonuclease activity was shown by the degradation of the primer after incubation with the DNA polymerase at 70°C for 1 h. As indicated in Fig. 8, the rate of *P. islandicum* DNA polymerase-associated 3'-5' exonuclease activity strongly depends on the pH of the reaction buffer. Using the optimal polymerase buffer (pH 7.3) the degradation of the primer by the enzyme from *P. islandicum* was significantly lower than with *Pwo* polymerase, whereas in buffer 2 (pH 8.6) the value for exonuclease activity was comparable to that of *Pwo* polymerase. Both on single-stranded primer and double-stranded primer/template the DNA polymerase from *P. islandicum* showed 3'-5' exonuclease activities in the absence of dNTPs. Even in the presence of 0.01 and 0.1 μ M dNTPs a *Taq* polymerase did not show any degradation of the primer, neither on single-stranded nor on double-stranded DNA substrate (Fig. 8). As expected for DNA polymerases with 3'-5' exonuclease activity, the degradation of the primer/template substrate by the DNA polymerases from *P. islandicum* and *P. woesei* was reduced in the presence of higher amounts (1 μ M) of dNTPs. Neither degradation nor elongation of the primer by the DNA polymerase from *P. islandicum* was detectable when the assay was performed at 37°C (data not shown). This result clearly indicates that the enzyme preparation did not contain a contaminating exonuclease activity from *E. coli* which could interfere with the test.

Applications

In this invention, we present the cloning procedure of a DNA polymerase gene and some biochemical properties of a new thermostable DNA polymerase from *Pyrobaculum islandicum*. The major uses of thermostable DNA polymerases are for in vitro amplification of DNA fragments and for determination of DNA sequences. *Taq* DNA polymerase is still used in many cases, but its fidelity is very low due to the absence of a 3'-5' exonuclease activity (Keohavong and Thilly 1989; Longley et al. 1990). Additionally, *Taq* polymerase has a thermostability of not more than few minutes at 100°C.

The DNA polymerase presented in this invention possesses a high thermostability at 90 and 100°C. It has to be investigated whether the presence of stabilizers such as nonionic detergents or bovine serum albumin (BSA) can even lead to an increased thermostability. The positive influence of the nonionic detergent octoxynol (commonly known as TRITON X-100) and BSA on the thermostability of DNA polymerases from *Pyrococcus furiosus* and *Thermococcus litoralis* has been described (EP Application no. 92118965 and 91303787.5). The thermostable DNA polymerase from *Pyrobaculum islandicum* is not irreversibly inactivated when subjected to the elevated temperatures for the time interval which is necessary to effect denaturation of double-stranded DNA, a key step in the

PCR process. As a result, the enzyme may provide additional advantages over the previously characterized thermostable enzymes. Higher temperatures up to 100°C may be required, e.g. as the GC composition of the DNA template is increased. For target sequences, denaturation temperatures over 95°C are needed. For these purpose, solvents such as glycerol or DMSO are included in PCR in order to partially destabilize the duplex DNA (Wong et al. 1991; Korge et al. 1992). These agents, in addition to destabilizing duplex DNA will affect primer stability, can inhibit activity and decrease thermostability of polymerases. In contrast, simply raising the denaturation temperature up to 95 or 100°C with *Pyrobaculum islandicum* DNA polymerase in an otherwise standard PCR can facilitate complete strand separation of PCR product eliminating the need for DNA helix destabilizing agents.

Furthermore, there is still a demand to obtain and produce a purified DNA polymerase having 3'-5' proofreading exonuclease activity with even high thermostability. The *P. islandicum* DNA polymerase with 3'-5' proofreading exonuclease activity has a substantially lower base incorporation error rate when compared with a non-proofreading DNA polymerase.

PCR performed with *P. islandicum* DNA polymerase. Thermostable DNA polymerases are particularly used for *in vitro* amplification of DNA fragments by PCR and for DNA sequencing. Due to the associated 3'-5' exonuclease activity, archaeal DNA polymerases offer the possibility to amplify DNA fragments with high fidelity (low mutation frequency). To date, all archaeal DNA polymerases used in PCR are derived from Euryarchaeota. No reports are available concerning the successful application of crenarchaeal DNA polymerases in PCR.

To verify the suitability of the DNA polymerases from *P. islandicum* in DNA amplification, PCR was performed using 1 U of enzyme and several primer combinations. As shown in Fig. 9, up to 1500 bp of a λ DNA template could be amplified. This is the first report on the possible use of a crenarchaeal DNA polymerase in PCR.

As above stated a method of a cloning procedure for B DNA polymerases using degenerated primers according to SEQ ID No. 1 and SEQ ID No. 2 is also subject of the present invention. Furthermore, a method of the amplification of fragments of a B DNA polymerase using primers according to SEQ ID No. 1 and 2 is subject of the present invention. The invention also includes the above mentioned methods if the following variants of the primers with SEQ ID No. 1 and 2 are used:

- primers wherein triplets are added or omitted, especially if 1-3 triplets are omitted and 1-5 triplets are added
- primers wherein one or more of the inosins shown in SEQ ID No. 1 or SEQ ID No. 2 are resubstituted by a naturally occurring base
- primers wherein one or more naturally occurring bases are substituted by inosin in addition to the substitutions according to SEQ ID No. 1 or SEQ ID No. 2, preferentially a naturally occurring base is substituted by inosin which is at the third position of a triplet.

Table 1: Influence of inhibitors on the DNA polymerase of *P. islandicum*
The influence of different inhibitors was detected at standard assay conditions (pH 7.5; 0.5 mM MgCl₂; 0 mM KCl; 75°C) using 0.5 U purified DNA polymerase. A preincubation of the enzyme with N-Ethylmaleimid was performed at 90°C for 20 min.

Inhibitor	Concentration	DNA polymerase activity
	mM	%
N-Ethylmaleimid	0	100
	1	40
	2	30
	5	0
	μM ($\mu\text{g/ml}$)	
Aphidicolin	0 (0)	100
	50 (17)	100
	200 (68)	70
	500 (170)	55
	2000 (680)	21
ddGTP	ddGTP/dGTP = 5	95
	10	80
	20	53

Example 1:

Strains, enzymes, vectors and chemicals

Pyrobaculum islandicum DSMZ4184 (Huber et al. 1987) was obtained from the Deutsche Sammlung von Mikroorganismen und Zellkulturen, GmbH (DSMZ; Braunschweig, Germany).

Magnesium cultures of *E. coli* DH5 α cells were included in the Expand Cloning Kit (Boehringer Mannheim). *E. coli* BL21 cells were obtained from Novagen Inc., Madison, USA.

Enzymes and kits for manipulations of DNA were obtained from Boehringer Mannheim or, if mentioned, from other customers. Kits for purification and gel extraction of DNA were products of QIAGEN (Hilden, Germany). A kit for preparation of plasmid and cosmid DNA was obtained from BioRad (München, Germany).

The expression vector pASK-IBA3, the inducer anhydrotetracycline and the Streptactin Sepharose columns were obtained from IBA, Göttingen, Germany.

Materials for the characterization of DNA polymerase were purchased from Boehringer Mannheim. N-Ethylmaleimid was obtained from Serva. Other chemicals for the preparation of media and buffers were obtained from Difco (Detroit, Wi), Serva (Heidelberg, Germany), Sigma (St. Louis, Mo.) and Merck (Darmstadt, Germany).

Isolation of DNA

P. islandicum was grown anaerobically at 98°C in 1.5 l Medium 390 as recommended by DMSZ. Cells were harvested, washed with TE buffer (10 mM Tris-HCl pH 8.0, 1 mM EDTA) and suspended in 3 ml 20% sucrose containing 10 mM Tris-HCl pH 8.0. Triton X-100 was added to a final concentration of 0.3%. After incubation at 20°C for 1 h, 3 ml of SET buffer (150 mM NaCl, 20 mM Tris-HCl pH 8.0, 1 mM EDTA), 0.32 ml of 10% sodium dodecyl sulfate and 0.16 ml of Proteinase K (10 mg/ml) were added. The mixture was incubated for 1 h at 37°C. After subsequent phenol-chloroform extraction, DNA was recovered by ethanol precipitation (Maniatis et al. 1982).

Example 2:

PCR and preparation of probes

For the first PCR, we designed two degenerated primer based on the conserved regions I and II of α -like DNA polymerases (Wang et al. 1989).

The forward primer was:

SEQ ID No. 1: 5'-GTI(CT)TIGATTT(CT)GCI(AT)(GC)I(CT)TITA(CT)CC-3', and

the reverse primer was:

SEQ ID No. 2: 5'-GAAIAGIGAGTCIGTATCICCAT-3',

using deoxy-inosin (I), which is complementary to all four nucleotides. Primers were custom synthesized by ARK Scientific, GmbH Biosystems (Darmstadt, Germany). Approx. one nanogram of *P. islandicum* DNA and 100 pmol of each primer were added to a PCR reaction mix containing 200 μ M deoxyribonucleoside triphosphates, the ExpandTM High Fidelity buffer system and 2 U of ExpandTM High Fidelity enzyme (Boehringer Mannheim). After an initial denaturation step at 94°C for 2 min, 30 cycles with a temperature profile of 10 sec at 94°C, 30 sec at 46°C and 90 sec at 68°C were performed with a DNA thermal cycler (GeneAmp PCR System 2400; Perkin-Elmer, Weiterstadt). Following the last cycle, the samples were incubated for additional 7 min at 68°C, to ensure completion of the extension step. The PCR product was purified and sequenced. Other PCR reactions were performed under the same conditions with different annealing temperatures and elongation times. The same PCR reaction mix, also containing the template and primer described above and 200 μ M dATP, dGTP and dCTP, but 190 μ M dTTP and 10 μ M DIG-11-dUTP (digoxigenin-labeled dUTP, alkali-stable; Boehringer Mannheim) was used to prepare a non-radioactive probe.

Genomic southern blot hybridization

P. islandicum DNA was digested over night with several restriction enzymes and resolved on an 0.7% agarose gel. The restricted DNA was transferred to a positively charged nylon membrane. The DIG-labeled probe described above was used to perform hybridizations at 68°C, and a three-step process, including membrane blocking, incubating with anti-DIG-antibody, Fab fragments and a reaction of the antibody-conjugated alkaline phosphatase with the chemiluminescent substrate CDP-StarTM (Boehringer Mannheim) was performed to detect the sites of the hybridized probe.

Cosmid cloning

Following the southern blot, the restriction enzyme *SacI* was chosen to perform a preparative digestion of *P. islandicum* DNA. Restricted DNA fragments with a size between 6 and 7.5 kbp were extracted from an agarose gel and cloned using the ExpandTM Cloning Kit (Boehringer Mannheim). The fragments were blunt-ended and ligated into the cosmid vector, and the constructs were packed into bacteriophages. An *E. coli* DH5 α magnesium culture was subsequently infected, and positive clones were selected by the vector-mediated ampicillin resistance. A cosmid clone containing the DNA polymerase gene was identified by colony filter hybridization (DIG System User's

Guide for Filter Hybridization), using the probe described. Cosmid DNA from a positive clone was prepared and sequenced.

DNA sequencing

Purified PCR products and cosmid DNA preparations were custom sequenced by SeqLab (Göttingen, Germany). The nucleotide sequences were determined by the dideoxy chain termination method (Sanger et al. 1977). All DNA samples were sequenced using ABI PRISM™ BigDye terminators, and analyzed by automated sequencers with a sequence analysis software (Applied Biosystems).

Example 3:

RNA isolation and determination of transcription start

In order to examine a putative transcription start region of the detected DNA polymerase gene, RNA from freshly cultivated *P. islandicum* cells was prepared, following a protocol written by DiRuggiero and Robb (1995), and a 5'RACE PCR was performed, using the 5'/3'RACE Kit (Boehringer Mannheim). For first-strand cDNA synthesis, cDNA amplification and control PCR, reverse primer deduced from the gene sequence were used. The amplified cDNA was gel extracted and directly sequenced.

Example 4:

Expression and purification of *P. islandicum* DNA polymerase

The DNA polymerase gene was expressed in *E. coli* using the vector pASK-IBA 3 (IBA, Göttingen). The expression cassette of this vector is transcriptionally controlled by the chemically inducible tetracycline promotor (Skerra 1994). For convenient purification of the recombinant protein, the vector contained a C-terminal, nine amino acids coding sequence, called Strep-tag. The Strep-tag allows to purify the resulting fusion protein in an affinity chromatography step on Sepharose-coupled streptavidin or Streptactin (Schmidt and Skerra 1994; Voss and Skerra 1997). Two primers, Exforw-*Bsa*I (5'-GCTGATGGTCTCGAATGGAATAAAAGTTTGGCCTCT-3') and Exrev-*Bsa*I (5'-GCTGATGGTCTCCGCGCTACTTAGAAAATCAAGAAGCGACCT-3') were used to amplify the polymerase gene from chromosomal *P. islandicum* DNA. The forward primer was designed such that a restriction site for *Bsa*I was generated and the start codon, coding for valin, was removed. The recombinant fusion protein started with the vector-coded methionin. The reverse primer was designed such that a restriction site for *Bsa*I was generated and the stop codon of the polymerase gene was removed. The PCR product was purified by gel extraction, cleaved with *Bsa*I, again gel extracted and ligated into *Bsa*I-restricted pASK-IBA 3 vector. Expression of the poly-

merase gene was performed in *E. coli* BL21. Transformed cells were cultivated at 37°C in 1000 ml of LB medium containing 100 µg ampicillin per ml. At an optical density of 0.5 at 600 nm, cells were induced by the addition of 100 µl anhydrotetracycline (2 mg/ml in dimethyl formamide; final concentration 200 µg/l), and the culture was incubated for a further 15 h. After cooling, the cells were harvested by centrifugation at 4°C, washed with buffer W, counteracting 50 mM Tris-HCl pH 8.0, 1 mM EDTA and the Complete mini, EDTA-free protease inhibitor cocktail (Boehringer Mannheim), and resuspended in 5 ml of buffer W. The cells were disrupted by sonication, and after centrifugation, the crude extract was heat treated at 80°C for 20 min and centrifuged again. The sample was applied to a Streptactin ready-to-use Sepharose column (1 ml), equilibrated with 5 ml buffer W. After washing the column twice with 5 ml buffer W, elution of the bound protein was effected by the stepwise application of 10 ml buffer W containing 2.5 mM desthiobiotin (Sigma). Active fractions were pooled and dialyzed against 500 volumes of 50 mM Tris-HCl pH 7.2 and 10% glycerol.

Example 5:

Polymerase activity assay

A non-radioactive colorimetric enzyme immunoassay (DNA Polymerase Assay, non-radioactive; Boehringer Mannheim) was performed, based on the incorporation of digoxigenin- and biotin-labeled dUTP, instead of radio-labeled nucleotides as used in the standard DNA polymerase assay, into the same DNA. As Suzuki et al. (1993) demonstrated, the results of a chemoluminescent enzyme-linked immunoassay, following the same principle as the test described here, displayed comparable results to the standard isotopic assay. The detection and quantification of the synthesized DNA followed the sandwich ELISA protocol, as recommended by the manufacture. Unless otherwise mentioned, the standard reaction mixture contained, in a 100 µl volume, 50 mM Tris-HCl pH 7.2, 0.5 mM MgCl₂, 60 µg BSA, 0.36 µM DIG-11-dUTP, 18 nM Biotin-11-dUTP, 18 µM each dNTP, 0.27 µg DNaseI-activated calf thymus DNA as a template, and the DNA polymerase sample. The reaction mix was incubated at 75°C for 1 h. *Taq* DNA polymerase was used as a standard, with 50 mM Tris-HCl pH 7.9, 5 mM MgCl₂ and 50 mM KCl in the reaction mix. One unit was defined as the amount of enzyme required to incorporate 10 nmol of dNTP into an acid-insoluble form at 75°C in 30 min. In order to determine the thermostability of the DNA polymerase from *P. islandicum*, preincubations were performed in a reaction mix without BSA, labeled or non-labeled dNTP and template-DNA, following by the addition of the missing components and standard incubation of 1 h at 75°C. For the determination of the biochemical properties of DNA polymerase, all reaction mixtures contained 0.5 U of enzyme.

Example 6:**DNA polymerase activity gel**

The detection of DNA polymerase activities on SDS polyacrylamid gels (SDS-PAGE) was performed according to a modified version (Niehaus et al. 1997) of the method described by Spanos and Hübscher (1983). Sample concentrations of one to five units of each DNA polymerase were run on SDS-PAGE (10%). Sample preparation and gel electrophoresis was carried out as described by Laemmli (1970). The running gel additionally contained 150 µg/ml DNaseI activated calf thymus DNA as template for the polymerases. DNA polymerases from *Thermus aquaticus* (*Taq* polymerase) and *Pyrococcus woesei* (*Pwo* polymerase) were purchased from Boehringer Mannheim. SDS was removed by immersing the gel 4 times in 50 mM Tris-HCl pH 7.2, 3 mM β-Mercaptoethanol, 1 mM EDTA, 5% glycerol. Renaturation of the proteins was performed at 4°C over night using 1 mM MgCl₂ and 1 µM dATP added to the same buffer. The gel was then incubated in activity buffer containing 50 mM Tris-HCl pH 7.2, 3 mM β-Mercaptoethanol, 1 mM DTT, 1 mM MgCl₂, 5% glycerol, 8 µM dGTP, dATP and dCTP, 4 µM dTTP and 5 µM DIG-11-dUTP, and incubated at 70°C for 4 h. DNA was transferred to a nylon membrane and incorporated DIG-11-dUTP was detected immunologically as described earlier.

Example 7:**Assay for exonuclease activity**

To analyze exonuclease activity, we incubated 1.5 U purified polymerase with 1 µg of a ³H-labeled DNA template, which was digested by sonication. The released ³H-labeled nucleotides were detected as cpm/10 min on a scintillizer after 4 h incubation. Incubations were carried out at 65°C and 37°C in the buffer (10 mM Tris-HCl, 1 mM DTT, 50 mM NaCl, 10 mM MgCl₂) normally used in this test.

Example 8:**Assay for 3'-5'-exonuclease activity on single strands**

For 3'-5' exonuclease activity on single-strands, the degradation of a synthetic primer (22-mer), 5'-labeled with digoxigenin for immunological detection, was analyzed. 3'-5' exonuclease activity on double-strands was measured using the same primer, hybridized to a template (34-mer). The sequences of primer and template are illustrated in Fig. 8. The assay principle was described previously (Niehaus, 1997 #77). 0.5 pmol of the labeled substrate and 0.1 U of purified DNA polymerase were incubated in 10 µl of different buffers without and with increasing amounts of dNTPs (up to 1 µM) at 70°C for 1 h. To ensure that no contaminating exonuclease activity from *E. coli* was

measured, the assay was further carried out at 37°C. The optimal DNA polymerase buffer containing 50 mM Tris-HCl pH 7.3 and 0.5 mM MgCl₂ was used as well as a buffer containing 50 mM Tris-HCl pH 8.6 and 0.5 mM MgCl₂. The reaction was stopped on ice with formamide buffer, and the samples were electrophoresed on a 12% polyacrylamide sequencing gel containing 8 M urea in TBE buffer. The degradation or polymerization products were visualized by immunological detection. *Pwo* polymerase in a buffer containing 10 mM Tris-HCl pH 8.9, 25 mM KCl, 2 mM MgSO₄ and 5 mM (NH₄)₂SO₄ and *Taq* polymerase in activity buffer described before were used as controls.

Example 9:

PCR reactions with *P. Islandicum*

PCR reactions with *P. islandicum* DNA polymerase were carried out with 1 U of enzyme in the following buffer: 15 mM Tris-HCl pH 8.6, 12.5 mM KCl, 2.5 mM (NH₄)₂SO₄, 1.25 mM MgCl₂, 20 µg/ml BSA. 10 ng of λ DNA was used as template. The forward primer 5'-GATGAGTTCGTGTCCGTACAACT-3' was combined with the following reverse primers: 5'-GGTTATCGAAATCAGCCACAGCG-3' for amplification of 500 bp, 5'-GTAACTTTGATTCTGGCCTGCG-3' for amplification of 1000 bp and 5'-GTGAGATAAACGGCAACTGCCGG-3' for amplification of 1500 bp. PCR cycles were carried out as described before with a temperature profile of 10 s at 94°C, 30 s at 56°C and 1.5 min at 75°C. Expand High Fidelity enzyme was used as a control under PCR conditions as recommended by the manufacturer.

Figure Legend

Fig. 1: Nucleotide sequence of the polymerase gene and deduced amino acid sequence of the family B DNA polymerase from *P. islandicum*

The nucleotides are numbered from the 5' site. The translation start site is at no. 131, the stop site at no. 2486.

Fig. 2: SDS Page (10%) of enzymatic fractions during purification of recombinant DNA polymerase from *P. islandicum*

Lanes: 1, crude extract; 2, crude extract after heat treatment (20'80°C); 3, molecular weight marker (broad range, BioRad); 4, enzyme preparation after chromatography on Streptactin. Molecular weights are indicated.

Fig. 3: DNA polymerase activity gel

Lanes: 1, *Taq* polymerase (5 U); 2, *P. islandicum* DNA polymerase enzyme preparation after chromatography on Streptactin (2 U); 3, crude extract from *P. islandicum*; 4, *Pwo* polymerase (2.5 U).

Fig. 4: Optimal pH for the DNA polymerase activity

The pH optimum was detected in a standard assay at 75°C with 0.5 mM MgCl₂.

Fig. 5: Optimal concentration of KCl, (NH₄)₂SO₄ and MgCl₂ for the DNA polymerase activity

The optimal concentration of MgCl₂ was measured in a standard assay at 75°C and pH 7.3. The standard assay was carried out with 0.2 U of purified DNA polymerase in the presence of various concentrations of KCl, (NH₄)₂SO₄ and MgCl₂.

Fig. 6: Influence of KCl on the DNA polymerase activity

The influence of monovalent potassium ions was detected in a standard assay at 75°C, pH 7.3 and with 0.5 mM MgCl₂.

Fig. 7: Thermostability of the recombinant DNA polymerase

To estimate the thermostability of the DNA polymerase, the enzyme was preincubated without BSA, nucleotides and template DNA at 90 and 100°C, following by the addition of all missing compounds and a standard incubation at 75°C, pH 7.3 and with 0.5 mM MgCl₂.

Fig. 8: Single- and double-strand dependent 3'-5' exonuclease activity of DNA polymerases from *Pyrobaculum islandicum* (*P. isl. pol*), *Thermus aquaticus* (*Taq pol*) and *Pyrococcus woesei* (*Pwo pol*). The assay was carried out, as described in Materials and Methods, at 70°C for 1 h with 0.1 U of each polymerase. A digoxigenin-labeled primer as a single-stranded substrate (ss) and a primer/template complex as a double-stranded substrate (ds) were used. Concentrations of dNTP (μM) in each reaction are indicated. The reaction products were separated on a 12% polyacrylamide gel containing 8 M urea and visualized by immunological detection. The sizes (nucleotides, nt) of the non-degraded primer and the reaction products are indicated. Lane P, digoxigenin-labeled primer in buffer without dNTPs; buffer 1:40 mM Tris-HCl pH 7.3, 0.5 mM MgCl₂; buffer 2:50 mM Tris-HCl pH 8.6, 0.5 mM MgCl₂. *Taq* and *Pwo* polymerase were tested using buffers according to Materials and Methods. Primer and template sequences are illustrated on the right.

Fig. 9: Application of DNA polymerase from *P. islandicum* in PCR. Reactions were carried out in a buffer containing 15 mM Tris-HCl pH 8.6, 12.5 mM KCl, 2.5 mM $(\text{NH}_4)_2\text{SO}_4$, 1.25 mM MgCl_2 and 20 $\mu\text{g/ml}$ BSA. 1 U of DNA polymerase from *P. islandicum* was used to amplify 500 bp (lane 1), 1000 bp (lane 2) and 1500 bp (lane 3) of a λ DNA template. Control PCR was performed with 2.5 U of High Fidelity enzyme (lane 4). 20 μl of each PCR product were applied on an 1% agarose gel. The corresponding molecular weights of the marker (lane 5) are indicated in kilobasepairs (kb).

References

- Blanco, L., Bernad, A., Blasco, M. A., & Salas, M. (1991) *Gene* 100, 27-38.
- Bradford, M. M. (1976) *Anal. Biochem.* 72, 248-254.
- Braithwaite, D. K., & Ito, J. (1993) *Nucleic Acids Res.* 21, 787-802.
- Cariello, N. F., Swenberg, J. A., & Skopek, T. R. (1991) *Nucleic Acids Res.* 19, 4193-4198.
- Datukishvili, N., Pokholok, D., Lottspeich, F., Prangishvili, D., & Rechinsky, V. (1996) *Gene* 177, 271-273.
- Dennis, P. P. (1997) *Cell* 89, 1007-1010.
- DiRuggiero, J., & Robb, F. T. (1995) in *Archaea-a laboratory manual. Thermophiles* (Robb, F. T., & Place, A. R., Eds.) pp 97-99, Cold Spring Harbor Laboratory Press.
- Dong, Q., Copeland, W. C., & Wang, T. S. (1993) *J. Biol. Chem.* 268, 24163-24174.
- Edgell, D. R., Klenk, H. P., & Doolittle, W. F. (1997) *J. Bacteriol.* 179, 2632-2640.
- Erlich, H. A., Gelfand, D. H., & Saiki, R. K. (1988) *Nature* 331, 461-462.
- Huber, B., Kristjansson, J. K., & Stetter, K. O. (1987) *Arch. Microbiol.* 149, 95-101.
- Keohavong, P., & Thilly, W. G. (1989) *Proc. Natl. Acad. Sci. USA* 86, 9253-9257.
- Korge, B. P., Gan, S. Q., McBride, O. W., Mischke, D., & Steinert, P. M. (1992) *Proc. Natl. Acad. Sci. USA* 89, 910-914.
- Kornberg, A., & Baker, D. (1992) *DNA replication*, Second ed., Freeman and company, New York.
- Laemmli, U. K. (1970) *Nature* 227, 680-685.
- Longley, M. J., Bennett, S. E., & Mosbaugh, D. W. (1990) *Nucleic Acids Res.* 18, 7317-7322.
- Lundberg, K. S., Shoemaker, D. D., Adams, M. W. W., Short, J. M., Sorge, J. A., & Marthur, E. J. (1991) *Gene* 108, 1-6.
- Maniatis, T., Fritsch, E. F., & Sambrook, J. (1982) *Molecular cloning: a laboratory manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
- Mullis, K., Faloona, F., Saiki, R., Horn, G., & Erlich, H. A. (1986) *Cold Spring Harbor Symp. Quant. Biol.* 51, 263-273.
- Niehaus, F., Frey, B., & Antranikian, G. (1997) *Gene* 204, 153-158.
- Perler, F. B., Kumar, S., & Kong, H. (1996) *Adv. Prot. Chem.* 48, 377-435.
- Pisani, F. M., Martino, C. D., & Rossi, M. (1992) *Nucleic Acids Res.* 20, 2711-2716.
- Prangishvili, D., & Klenk, H. P. (1993) *Nucleic Acids Res.* 21, 2768.
- Sanger, F., Nicklen, S., & Coulson, A. R. (1977) *Proc. Natl. Acad. Sci. USA* 74, 5463-5467.
- Schleper, C., DeLong, E. F., Preston, C. M., Feldman, R. A., Wu, K.-Y., & Swanson, R. V. (1998) *J. Bacteriol.* 180, 5003-5009.
- Schleper, C., Swanson, R. V., Mathur, E. J., & DeLong, E. F. (1997) *J. Bacteriol.* 179, 7803-7811.

- Schmidt, T. G. M., & Skerra, A. (1994) *J. Chromatogr. A* 676, 337-345.
- Skerra, A. (1994) *Gene* 151, 131-135.
- Southworth, M. W., Kong, H., Kucera, R. B., Ware, J., Jannasch, H. W., & Perler, F. B. (1996) *Proc. Natl. Acad. Sci. USA* 93, 5281-5285.
- Spanos, A., & Hübscher, U. (1983) *Meth. Enzymol.* 91, 263-277.
- Stetter, K. O. (1996) *FEMS Microbiol. Rev.* 18, 149-158.
- Suzuki, K., Craddock, B. P., Kano, T., & Steigbigel, R. T. (1993) *Anal. Biochem.* 210, 277-281.
- Uemori, T., Ishino, Y., Doi, H., & Kato, I. (1995) *J. Bacteriol.* 177, 2164-2177.
- Voss, S., & Skerra, A. (1997) *Protein Eng.* 10, 975-982.
- Wang, T. S. F., Wong, S. W., & Korn, D. (1989) *FASEB J.* 3, 14-21.
- Woese, C. R., & Fox, G. E. (1977) *Proc. Natl. Acad. Sci. USA* 74, 5088-5090.
- Wong, D. M., Weinstock, P. H., & Wetmur, J. G. (1991) *Nucleic Acids Res.* 19, 2251-2259.

Claims

1. A DNA polymerase obtainable from *Pyrobaculum islandicum* whereas six conserved motifs indicative of family B DNA polymerases were contained in the *Pyrobaculum islandicum* DNA polymerase sequence and whereas this polymerase exhibits 3'-5'-exonuclease activity.
 2. A DNA polymerase according to claim 1 whereas the molecular weight of this polymerase is approximately 90 kDa.
 3. A DNA polymerase according to claims 1 or 2 whereas the polymerase exhibits most activity at pH 7.3.
 4. A DNA polymerase according to one of the claim 1-3 whereas this polymerase exhibits a low sensitivity to ddGTP.
 5. A recombinant DNA polymerase according to one of the chains 1-4 obtainable from E.coli BL21 transformed by a vector containing a gene which encodes the DNA polymerase.
 6. DNA encoding a polymerase according to one of the claims 1-5.
 7. Vector containing a gene of a polymerase according to one of the claims 1-5.
 8. Host transformed by a vector according to claim 7.
 9. Method for producing a recombinant DNA polymerase according to claim 5.
 10. Use of a DNA polymerase according to one of the claims 1-5 for in vitro amplification of DNA fragments.
 11. Use of a DNA polymerase according to one of the claims 1-5 for determination of DNA sequences.
 12. A method for cloning a B DNA polymerase using primers with SEQ ID No. 1 and SEQ ID No. 2 or variants thereof.
-

13. A method of the amplification of fragments of B DNA polymerase using primers with SEQ ID No. 1 and SEQ ID No. 2 or variants thereof.

Fig. 1/1

SEQ ID No. 3

SEQ ID No. 4

GCTTTTAAAGTAGCCACTCGATTACATAGTTGTGCGCCGTGCCCGGTTTTCGAGGTAATTGTTTAAAGCATTCCGCCACAGCGTTCATAT 90
ACTGTAAATCAATTTATTTTATATTGTTTCAAGCATTGTGGAATAAAAGTTTGGCCTCTCGACATTACATATGCCGTAGTTGGTAGT 180
V E L K V W P L D I T Y A V V G S
GTTCTGAGATTAGGATTTTCGGTATTTAAGCAGCGGTGAGCGTGTGTACTTATCGATAGATCTTTAAGCCGTATTTCTACGTCGAT 270
V P E I R I F G I L S S G E R V V L I D R S F K P Y F Y V D
TGTGCCGTTTGTGAACCAGCTGCTCTAAAGACTGCGTTGTGCGGAGTTGCGCCAATAGACGATGTACAAATAGTTGAAAGGAGGTTCTCTC 360
C A V C E P A A L K T A L S R V A P I D D V Q I V E R R F L
GGAAGGTCTAAGAAGTTTGTAAAGTTATTGCTAAATTCCTGAGGATGTCAGGAACTTAGAGAAGCCGCAATGTCGATACCAAGGGTT 450
G R S K K F L K V I A K I P E D V R K L R E A A M S I P R V
TCTGGCGTTTATGAAGCAGATATTCGTTTTACATGCGATATATGATAGACATGGGGGTAGTGCCTGTAGTTGGAATGTGGCAGAGGTA 540
S G V Y E A D I R F Y M R Y M I D M G V V P C S W N V A E V
GAAGAGGGGGGTAGACTTGGCGGGATTCCACATACGTTGTCTCTCAGTGGTACGGGATTGACGAAGGCTTTCCCTTTCATTGAAAGTT 630
E E G G R L G G I P T Y V V S Q W Y G I D E G F P P S L K V
ATGGCGTTTGATATCGAGGTTTATAACGAGCGCGGCTCTCCCGATCCTATCAGAGATCCCGTGGTTATGTTAGCCATAAAGACAAACGAT 720
M A F D I E V Y N E R G S P D P I R D P V V M L A I K T N D
GGGCACGAGGAGGTTTTTGAAGCTAGTGGGAAAGACGATAGGGGGGTTGTGCGCGCTTTTGTGGATTTTATCAGAAGTTATGACCCCGAT 810
G H E E V F E A S G K D D R G V V R A F V D F I R S Y D P D
GTAATTGTTGGCTATAATTCCAACGCGCTTTCGATTGCGCTTATTTAGTAGAGCGTGCGAAAGCCGTGCGCGTGCCTCAAGGTTGATAGA 900
V I V G Y N S N G F D W P Y L V E R A K A V G V P L K V D R
CTGAGCAATCCCTCAACAGAGCGTGTATGGACACTGGTCTATCGTTGGTAGAGCAAATGTAGATCTCTATAACATCGTGGAAAGAGTTTC 990
L S N P P Q Q S V Y G H W S I V G R A N V D L Y N I V E E F
CCGGAGATTAAGCTAAAGACTCTTGACCGTGTGCTGAATATTTGGTGTAAATGAAAAGAGAAGAGAGGGTGTGATCCCAGGCCACAAG 1080
P E I K L K T L D R V A E Y F G V M K R E E R V L I P G H K
ATTTATGAATACTGGAAAGACCCCAATAAAAGACCTCTACTTAAGCGGTATGTTCTAGACGACGTACGATCCACTCTAGGACTTGCCGAC 1170
I Y E Y W K D P N K R P L L K R Y V L D D V R S T L G L A D
AAGCTCCTACCGTTTTTAATACAACGTCTTCTGTATCTGGGCTACCGCTGGATCAGGTGGCAGCGCGAGCGTGGGCAACAGGGTAGAG 1260
K L L P F L I Q L S S V S G L P L D Q V A A A S V G N R V E
TGGATGCTTCTTAGGTACGCGTACCGCTGGCGAGGTAGCTCCGAACAGAGAGGAGGGAGTACGAACCATATAAGGGGGCCATCGTG 1350
W M L L R Y A Y R L G E V A P N R E E R E Y E P Y K G A I V
CTTGAGCCAAAGCCCGGATGTACGAAGACGTGCTTGTACTTGACTTCTCTTCCATGTACCCTAACATCATGTAAGTACAACCTATCG 1440
L E P K P G M Y E D V L V L D F S S M Y P N I M M K Y N L S
CCGGATACGTACCTAGAGCCCGCGAGCGGATCCGCCCGAGGGTGTAACGTGGCCCCGAGGTGGGGCATAGGTTTAGGAGAAGCCCA 1530
P D T Y L E P G E P D P P E G V N V A P E V G H R F R R S P

Fig. 1/2

CCAGGCTTCGTCCTCAGGTGTTGAAGAGCTTGGTGGAGCTTAGAAAGGCGGTAAGAGAGGAGGCGAAGAAATACCCCCAGATTGCGCA 1620
P G F V P Q V L K S L V E L R K A V R E E A K K Y P P D S P

GAGTTTAAGATCTTAGACGAGAGACAACGTGCCCTCAAGGTTATGGCCAATGCCATTTATGGATACTTGGGCTGGGTGGGGCCCCGGTGG 1710
E F K I L D E R Q R A L K V M A N A I Y G Y L G W V G A R W

TATAAGCGGGAGGTAGCCGAGTCTGTGACGGCTTTTGCCAGAGCGATTCTAAAAGACGTTATTGAACAAGCTAGAAGGTTGGGCATTGTG 1800
Y K R E V A E S V T A F A R A I L K D V I E Q A R R L G I V

GTGTATATGGCGATACAGACAGCCTATTTGTCAAAAAACATGGAGACGTGGACAACTGATCAAGTACGTGGAGGAGAAGTACGGCATA 1890
V V Y G D T D S L F V K K H G D V D K L I K Y V E E K Y G I

GACATAAAGGTGGACAAGGATTACGCCAAGGTGCTTTTCACGGAGGCTAAGAAGAGGTACGCTGGCTTATTGAGAGATGGCGTATAGAT 1980
D I K V D K D Y A K V L F T E A K K R Y A G L L R D G R I D

ATTGTGGGATTTCGAGGTCTGTAGGGGAGACTGGAGTGAACCTTGCCAAAGACGTTTCAGCTAAGAGTTATAGAGATTATACTTAAGTCAAGA 2070
I V G F E V V R G D W S E L A K D V Q L R V I E I I L K S R

GATATTGTGGAAGCTAGACATGGCGTTATAAAATATATTAGAGAAATAATAGAACGACTGAAAAATTACAAGTTTAACATTGATGATTTA 2160
D I V E A R H G V I K Y I R E I I E R L K N Y K F N I D D L

ATAATATGAAAACTCTAGACAAAGAGTTAGACGAATATAAGGCGTATCCGCCTCATGTTACAGCTGCGCAAATTCTCAAGAGACACGGC 2250
I I W K T L D K E L D E Y K A Y P P H V H A A Q I L K R H G

TATCGAGTGGGCAAAGGCACTACGATAGGCTATGTCATAGTTAAAGGAGGCGAAAAAGTATCTGAGAGGGCGTTGCCGTATATACTCCTT 2340
Y R V G K G T T I G Y V I V K G G E K V S E R A L P Y I L L

GACGATATAAAAAAGATAGATATCGACTACTACATAGAGAGACAGATAATACCAGCGGCGTTGAGAATAGCAGAGGTTATAGGGGTGAAA 2430
D D I K K I D I D Y Y I E R Q I I P A A L R I A E V I G V K

GAGTCGGATCTTAAACGGGGAGAATGAAAGGTGCTTCTTGATTTTCTAAGTTAGAAAACTACGTTTCTGCGTTATTGATCAAAAG 2520
E S D L K T G R M E R S L L D F L S .

TCATATCTTCATATCTTCAGGCACCTTTTGTAGAGGGCCAACTCTAGCAGTTATAATGACTTTGACTCCCCTTTTCGTGGCTAAGTCCA 2610




Figure 2





Figure 3

Fig. 4






Fig. 5

Fig. 6




Fig. 7





Figure 8






Figure 9

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 00/01917

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/54 C12N9/12 C12Q1/68

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 624 641 A (HOFFMANN LA ROCHE) 17 November 1994 (1994-11-17) the whole document ---	
A	WO 93 25691 A (CONSIGLIO NAZIONALE RICERCHE ;PISANI FRANCESCA MARIA (IT); ROSSI M) 23 December 1993 (1993-12-23) the whole document ---	
A	WO 99 07837 A (DIVERSA INC) 18 February 1999 (1999-02-18) page 33, line 11 - line 30 --- -/--	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

G document member of the same patent family

Date of the actual completion of the international search

9 June 2000

Date of mailing of the international search report

28/06/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Hornig, H

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 00/01917

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>BURGGRAF S ET AL: "An intron within the 16S ribosomal RNA gene of the archaeon <i>Pyrobaculum aerophilum</i>."</p> <p>PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED STATES OF AMERICA 1993, vol. 90, no. 6, 1993, pages 2547-2550, XP002139882</p> <p>ISSN: 0027-8424</p> <p>the whole document</p> <p>---</p>	
A	<p>M. MOLITOR ET AL.: "A dissimilatory sirohaem-sulfite-reductase-type protein from the hyperthermophilic archaeon <i>Pyrobaculum islandicum</i>"</p> <p>MICROBIOLOGY, vol. 144, 1998, pages 529-541, XP002139883</p> <p>READING, UK</p> <p>the whole document</p> <p>---</p>	
A	<p>EDGEALL DAVID R ET AL: "Gene duplications in evolution of archaeal family B DNA polymerases."</p> <p>JOURNAL OF BACTERIOLOGY 1997, vol. 179, no. 8, 1997, pages 2632-2640, XP002139884</p> <p>ISSN: 0021-9193</p> <p>cited in the application</p> <p>---</p>	
A	<p>SCHLEPER CHRISTA ET AL: "Characterization of a DNA polymerase from the uncultivated psychrophilic archaeon <i>Cenarchaeum symbiosum</i>."</p> <p>JOURNAL OF BACTERIOLOGY DEC., 1997, vol. 179, no. 24, December 1997 (1997-12), pages 7803-7811, XP002139885</p> <p>ISSN: 0021-9193</p> <p>cited in the application</p> <p>the whole document</p> <p>---</p>	
A	<p>C. SCHLEPER ET AL.: "Genomic analysis reveals chromosomal variation in natural populations of the uncultured psychrophilic archaeon <i>Cenarchaeum symbiosum</i>"</p> <p>J. BACTERIOL., vol. 180, no. 19, October 1998 (1998-10), pages 5003-5009, XP002139886</p> <p>AM. SOC. MICROBIOL., BALTIMORE, US;</p> <p>cited in the application</p> <p>the whole document</p> <p>---</p> <p style="text-align: center;">-/--</p>	

INTERNATIONAL SEARCH REPORT

Intern. Appl. No.

PCT/EP 00/01917

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>BARNS SUSAN M ET AL: "Remarkable archaeal diversity detected in a Yellowstone National Park hot spring environment." PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED STATES OF AMERICA 1994, vol. 91, no. 5, 1994, pages 1609-1613, XP002139887 ISSN: 0027-8424 the whole document</p> <p>---</p>	
A	<p>J. KJEMS ET AL.: "Phylogenetic relationship amongst the hyperthermophilic archaea determined from partial 23S rRNA gene sequences" SYSTEM. APPL. MICROBIOL, vol. 15, no. 2, 1992, pages 203-208, XP000914637 FISCHER, STUTTGART, FRG the whole document</p> <p>---</p>	
A	<p>HUBER R ET AL: "PYROBACULUM NEW-GENUS OF NEUTROPHILIC ROD-SHAPED ARCHAEABACTERIA FROM CONTINENTAL SOLFATARAS GROWING OPTIMALLY AT 100 C" ARCHIVES OF MICROBIOLOGY 1987, vol. 149, no. 2, 1987, pages 95-101, XP000914645 ISSN: 0302-8933 the whole document</p> <p>---</p>	
P,X	<p>M. KAEHLER AND G. ANTRANAKIAN : "Pyrobaculum islandicum family B DNA polymerase (polB3) gene, complete cds." EMBL SEQUENCE DATABASE, 26 January 2000 (2000-01-26), XP002139889 Hinxton, UK Accession no. AF195019;</p> <p>---</p>	1-9
P,X	<p>DATABASE BIOSIS 'Online! BIOSCIENCES INFORMATION SERVICE, PHILADELPHIA, PA, US February 2000 (2000-02) KAEHLER MARKUS ET AL: "Cloning and characterization of a family B DNA polymerase from the hyperthermophilic crenarchaeon Pyrobaculum islandicum." Database accession no. PREV200000113650 XP002139890 abstract & JOURNAL OF BACTERIOLOGY FEB., 2000, vol. 182, no. 3, February 2000 (2000-02), pages 655-663, ISSN: 0021-9193</p> <p>-----</p>	1-13

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No
PCT/EP 00/01917

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0624641 A	17-11-1994	US 5491086 A	13-02-1996
		CA 2123354 A	15-11-1994
		JP 6343466 A	20-12-1994
WO 9325691 A	23-12-1993	IT 1255666 B	09-11-1995
		AU 4512693 A	04-01-1994
WO 9907837 A	18-02-1999	US 5948666 A	07-09-1999
		AU 8914398 A	01-03-1999
		EP 1002058 A	24-05-2000

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)